

Nishimori point in the 2D $\pm J$ random-bond Ising model

A. Honecker¹, M. Picco² and P. Pujol³

¹*Institut für Theoretische Physik, TU Braunschweig, Mendelssohnstr. 3, 38106 Braunschweig, Germany.*

²*LPTHE*, Université Pierre et Marie Curie, Paris VI, Université Denis Diderot, Paris VII
Boite 126, Tour 16, 1^{er} étage, 4 place Jussieu, 75252 Paris Cédex 05, France.*

³*Laboratoire de Physique*, Groupe de Physique Théorique, ENS Lyon, 46 Allée d'Italie, 69364 Lyon Cédex 07, France.
(October 9, 2000; revised April 24, 2001)*

We study the universality class of the Nishimori point in the 2D $\pm J$ random-bond Ising model by means of the numerical transfer-matrix method. Using the domain-wall free-energy, we locate the position of the fixed point along the Nishimori line at the critical concentration value $p_c = 0.1094 \pm 0.0002$ and estimate $\nu = 1.33 \pm 0.03$. Then, we obtain the exponents for the moments of the spin-spin correlation functions as well as the value for the central charge $c = 0.464 \pm 0.004$. The main qualitative result is the fact that percolation is now excluded as a candidate for describing the universality class of this fixed point.

PACS numbers: 75.50.Lk, 05.50.+q, 64.60.Fr

In the past years, the subject of disordered systems has known a huge renewal of interest in the condensed matter and statistical mechanics community. Among these disordered models, two-dimensional systems are of a particular interest. Since the discovery of the unitary series of conformal field theory (CFT) in 1984 [1], exact values for the exponents of many well known models of statistical mechanics have been given. However, an equivalent classification for universality classes of such systems in the presence of impurities is still missing. A first big step towards a more general classification has been done recently as a random matrices classification [2]. The data for critical exponents in most of the experimental relevant fixed points for impure system is however still not available.

The Ising model on a square lattice is one of the most popular two-dimensional systems. It is specified by the energy of a spin configuration

$$E(\{S_i\}) = \sum_{\langle i,j \rangle} J_{i,j} \delta S_i S_j, \quad (1)$$

where the sum is over all bonds and the coupling constants $J_{i,j}$ are bond dependent. We consider here the $J_{i,j} = \pm 1$ Random-Bond Ising Model (RBIM) with the following probability distribution:

$$P(J_{i,j}) = p\delta(J_{i,j} - 1) + (1 - p)\delta(J_{i,j} + 1). \quad (2)$$

Note that with these conventions, the pure model ($p = 0$) is characterized by $J_{i,j} = -1$ and thus has a ferromagnetic groundstate.

The RBIM is similar to other relevant disordered models such as the Chalker-Coddington random network model which was proposed originally in the context of the quantum Hall effect plateau transition [3]. However, it is important to stress that these systems have a different phase diagram and therefore their fixed points have no reason to be in the same universality class [4].

The topology of the phase diagram of the RBIM depends crucially on the type of disorder one considers. An

instructive example is provided by a disorder having only two possible values for the bonds with equal signs and probabilities. It is by now well established [5] that the only non-trivial fixed points are located at the extrema of the boundary of the ferromagnetic phase, corresponding to the pure Ising fixed point and a zero temperature fixed point which turns out to be in the percolation universality class. It is interesting to notice that percolation is also the universality class of the so-called spin quantum Hall model [6], another random network model.




FIG. 1. Phase diagram of the two-dimensional $\pm J$ random-bond Ising model.

When the distribution contains also bonds with different signs (like in (2)), the situation is more subtle. For a certain class of probability distributions, Nishimori has shown that a so-called ‘Nishimori’ line exists where many properties can be calculated exactly [7]. For the probability distribution (2), this line is given by

$$e^\beta = \frac{1 - p}{p}, \quad (3)$$

with $\beta = 1/T$. On the Nishimori line, the internal energy can be calculated exactly and an upper bound can be given for the specific heat. Also of interest is an equality of the moments of the spin correlation functions (see below). Nishimori has further proven inequalities for the correlation functions which yield important constraints on the topology of the phase diagram which is shown in Fig. 1 for the $\pm J$ RBIM [8]. Since the Nishimori line is also invariant under Renormalization Group (RG) transformations [10], the intersection of the Nishimori line and the Ferro-Para transition line must be a fixed point. This so-called Nishimori point (N) corresponds to a new universality class belonging precisely to the family of strong disorder fixed points. The bold line in the phase diagram Fig. 1 is the phase boundary between the ferromagnetic and paramagnetic regions. At zero temperature the model has a spin glass phase [11]. The three non-trivial fixed points along the bold line are the pure Ising fixed point, the Nishimori point at the crossing with the Nishimori line (dotted line) and the zero temperature point, separating the ferro and the spin glass phases. The properties of the latter point are still mostly unknown. This point will be the subject of future investigations [12].

In the last years, many numerical and analytical efforts have been made in order to identify the universality class of the Nishimori point. Very recently, an analytic approach suggested that it is governed by an $Osp(2n+1|2n)$ symmetry (and maybe $Osp(2n+2|2n)$ [9], but unfortunately, the classification of CFTs with such symmetries is still missing. From a numerical point of view, there is by now a long list of results [13–17]. An important observation is that all the numerical results for the critical exponents tend to suggest that this point is in the percolation universality class. Because of the importance of the statistical model on its own and its relevance for understanding the plateau transition in the quantum Hall effect, it is crucial to elucidate the similarity to percolation and the relation to the super-symmetric CFT proposed in the literature.

In this letter we provide results of extensive numerical transfer-matrix calculations of the Nishimori point with the binary distribution (2) for bonds on the square lattice. We use the domain-wall free-energy to accurately locate the critical concentration of disorder p_c and to estimate the exponent ν . Then, we analyze the spin correlation functions and the scaling of the free energy, giving accurate and novel results for the magnetic exponent η and central charge c . Apart from improving the identification of this universality class (providing in particular values for the central charge which have never been measured before), our main result is that percolation is excluded as a possible candidate for describing this fixed point.

First, we use the free energy of a domain wall [13] to locate the critical point. For a strip of width L the domain-wall free-energy d_L is defined as [18]

$$d_L = L^2 \left(f_L^{(p)} - f_L^{(a)} \right), \quad (4)$$

where $f_L^{(p)}$ is the free energy *per site* of a strip of width L with *periodic* boundary conditions and $f_L^{(a)}$ the corresponding one with *antiperiodic* boundary conditions. d_L is an observable which can be used directly to study the RG flow under scale transformations. In particular, it is constant at a fixed point.

We have computed $f_L^{(p)} = \frac{\ln Z^{(p)}}{LN}$ and $f_L^{(a)} = \frac{\ln Z^{(a)}}{LN}$ employing a standard transfer matrix technique with sparse matrix factorization (see, e.g., [19]) on strips of length $N = 10^6$. Since randomness is strong, care must be taken to reduce fluctuations even if the free energies are self-averaging. Therefore, we have fixed the concentration of bonds p globally on a sample and computed $f_L^{(p)}$ and $f_L^{(a)}$ on the same sample. Still, one needs around 1000 to 4000 samples of $L \times 10^6$ strips to obtain sufficiently small error bars for $L \leq 12$. Even on modern computers this needs an amount of CPU time which precludes the analysis of wider strips. However, since we are looking for a fixed point, no crossover effects are expected and it is legitimate to use small system sizes.




FIG. 2. Domain-wall free-energy. The inset shows the raw data and the main panel the scaling collapse with $p_c = 0.1094$ and $\nu = 1.33$. The symbols are for $L = 8$ (filled boxes), $L = 9$ (open boxes), $L = 10$ (*), $L = 11$ (x) and $L = 12$ (+). Error bars are much smaller than the size of symbols.

The inset of Fig. 2 shows $d_L(p)$ along the Nishimori line (3) in the vicinity of the critical concentration p_c . A finite-size estimate for p_c is given by the crossing points $d_{L_1}(p_c) = d_{L_2}(p_c)$. After extrapolation to an infinitely wide strip (details will be given elsewhere [12]), one obtains

$$p_c = 0.1094 \pm 0.0002. \quad (5)$$

This estimate improves upon the accuracy of earlier estimates [14–17]. It agrees perfectly with the transfer matrix computations [14,17] while we find a slightly smaller

value of p_c than [15,16]. We would like to mention that (5) is confirmed by standard Monte Carlo simulations on systems up to 32×32 sites [12] – the present estimate is just more accurate.

One can extract also the correlation length exponent ν from d_L if one assumes the scaling form

$$d_L(p - p_c) = d \left((p - p_c) L^{1/\nu} \right). \quad (6)$$

Again, we omit details [12] and quote just the final result

$$\nu = 1.33 \pm 0.03. \quad (7)$$

The main panel of Fig. 2 demonstrates that d_L follows indeed the scaling form (6) with the parameters (5) and (7).

The result (7) is in complete agreement with $\nu = 1.32 \pm 0.08$ obtained by high-temperature series [15] as well as the value $\nu = 4/3$ for percolation (see, e.g., [20]).

Another important quantity is the magnetic exponent η . This exponent can be measured, for example, by computing spin-spin correlation functions. As we mentioned in the introduction, all along the Nishimori line the moments of these correlation functions are equal two by two:

$$[\langle S(x_1, y_1) S(x_2, y_2) \rangle^{2k-1}] = [\langle S(x_1, y_1) S(x_2, y_2) \rangle^{2k}] \quad (8)$$

for any integer k . Here $[\dots]$ stands for the average over the disorder. Assume now that the correlation functions (8) decay algebraically on a plane and define by x, y the coordinates on the infinite cylinder of circumference L , with $x \in [1, L]$ and $y \in]-\infty, +\infty[$. Using a conformal mapping, one infers then the following behavior of the correlation functions on the cylinder:

$$[\langle S(x_1, y) S(x_2, y) \rangle^n] \propto \left(\sin \left(\frac{\pi(x_2 - x_1)}{L} \right) L \right)^{-\eta_n}. \quad (9)$$

For a pure system, one has $\eta_n = n \times \eta$. On the other hand, in the case of percolation over Ising clusters, it is easy to see that the moments of spin correlation functions are all equal (and not only two by two). Then, if the Nishimori point is in the percolation universality class, the exponents for the correlation functions in (9) should collapse to a unique value $\eta_n = \eta$ at the critical point.

In order to verify this we have calculated the spin-spin correlation functions on cylinders of width L and length $400 \times L$, (*i.e.* with the length $\gg L$) for L up to 20. We have checked that for width $L = 12$, lattice and finite length corrections are of order 1%. One example of these correlation functions can be seen in Fig. 3 (with $x_1 = y = 0$ and $x_2 = x$) on a doubly logarithmic scale. One observes that the correlation functions nicely obey the power law (9), thus verifying both the correct location of the critical point as well as the functional form of the spin-spin correlation function in a finite strip.




FIG. 3. Moments of the spin-spin correlation function for $p = 0.1095$ and $L = 20$. We only show the odd moments: $n = 1$ (+), $n = 3$ (×), $n = 5$ (*) and $n = 7$ (open boxes). Error bars are smaller than the size of the symbols. The values of the exponents are given in (10).

We can then fit the exponent by studying the dependence with distance of the correlation functions (9) or by studying the dependence with L for the fixed location $x = L/2$. The first method has proven to give smaller error bars and we obtain for the family of exponents η_n for $p_c = 0.1095$ and $L = 20$:

$$\begin{aligned} \eta_1 &= \eta_2 = 0.1854 \\ \eta_3 &= \eta_4 = 0.2561 \\ \eta_5 &= \eta_6 = 0.3015 \\ \eta_7 &= \eta_8 = 0.3354, \end{aligned} \quad (10)$$

with relative errors at most of the order of 1%.

One immediately notices two things:

- i) The value for η_1 differs considerably from the value of percolation $\eta = 5/24 \approx 0.2083$ (see, e.g., [20]),
- ii) the exponents for higher moments are also considerably different from η_1 which is also clear from inspection of Fig. 3.

These results are compatible with the behavior of the magnetic susceptibility that will be presented elsewhere [12]. We have also calculated estimates for the exponents assuming different values for p_c , namely, $p_c = 0.109$ and 0.110 and the results are still distinct from the ones of percolation (we obtain $\eta_1 = 0.180(1)$ for $p = 0.109$ and $\eta_1 = 0.190(1)$ for $p = 0.110$). Moreover, it is only in the region very close to $p = 0.1095$ that we obtain a stable estimate for η_1 as we increase the width L of the lattices. One can then conclude from the exponents controlling the algebraic decay of the correlation functions that the Nishimori point is not in the percolation universality class.

Finally, we discuss the (effective) central charge c . It characterizes the number of gapless degrees of freedom at the critical point and appears as the universal coefficient

of the first finite-size correction to the free energy for periodic boundary conditions [21]

$$f_L^{(p)} = f_\infty^{(p)} + \frac{c\pi}{6L^2} + \dots \quad (11)$$

The leading term $f_\infty^{(p)}$ is not universal and does indeed change already when we modify the conventions for the model (1). There are higher-order finite-size corrections to the free energy including terms of the form L^{-4} . The central charge is an important quantity identifying the CFT description [1] of a fixed point. One has $c = 1/2$ for the critical point of the pure Ising model, but it has not been determined yet for the Nishimori point.

In the process of computing d_L we have also obtained estimates of $f_L^{(p)}$ for different values of p . One can either fit these values for $f_L^{(p)}$ exactly by (11) ignoring further corrections in which case the data for the smallest values of L should not be used. Or one includes a correction term of the form L^{-4} which improves the convergence with system size. These two approaches yield consistent estimates for a given p . In addition, one can test that the result does not change significantly if other higher-order corrections are added. It should also be noted that the sensitivity of the estimates for c with respect to the location of p_c is negligible in comparison with the errors coming from the finite-size analysis. The final result is that the following is a safe estimate for c at the Nishimori point of the $\pm J$ RBIM (more details will be given in [12]):

$$c = 0.464 \pm 0.004. \quad (12)$$

Assuming again the universality class of percolation, we would expect the value for percolation in the Ising model $c = \frac{5\sqrt{3}\ln 2}{4\pi} \approx 0.4777$ [5]. Even if our result (12) is close to this value, it can still be distinguished safely from percolation. This finding is one more argument that the Nishimori point is *not* in the universality class of percolation, at least the one expected from Ising clusters. Notice also that the central charge does not rely on a choice of observables, and this argument can then be considered as the most general one.

The results presented in this letter provide new insight into the 2D $\pm J$ RBIM and the Nishimori point. The main result is that, according to the magnetic exponent and the behavior of the higher moments of the correlation functions, the universality class is different from the one of percolation, at least considering the Ising spin variables as fundamental observables. This conclusion is supported by the central charge which is independent of the choice of observables and was measured here for the first time. These results and a detailed study of the zero temperature fixed point will provide a general and complete control of the RBIM, which one could consider as the simplest, but most fundamental model for disordered systems in 2D.

We would like to thank S. Franz, J. L. Jacobsen, J. M. Maillard, G. Mussardo, N. Read and F. Ritort for useful discussions and comments.

[◊] Unité Mixte de Recherche CNRS UMR 7589.

^{*} URA 1325 du CNRS associée à l'Ecole Normale Supérieure de Lyon.

- [1] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, *Nucl. Phys. B* **241**, 333 (1984), *J. Statist. Phys.* **34**, 763 (1984).
- [2] M. R. Zirnbauer, *J. Math. Phys.* **37**, 4986 (1996), A. Altland and M. R. Zirnbauer, *Phys. Rev. B* **55**, 1142 (1997).
- [3] S. Cho and P. A. Fisher, *Phys. Rev. B* **55**, 1025 (1997).
- [4] J. T. Chalker, N. Read, V. Kagalovsky, B. Horovitz, Y. Avishai and A. W. W. Ludwig, *cond-mat/0009463*.
- [5] J. L. Jacobsen and J. L. Cardy, *Nucl. Phys. B* **515**, 701 (1998).
- [6] I. A. Gruzberg, N. Read and A. W. W. Ludwig, *Phys. Rev. Lett.* **82**, 4524 (1999), J. Cardy, *Phys. Rev. Lett.* **84**, 3507 (2000).
- [7] H. Nishimori, *Prog. Theor. Phys.* **66**, 1169 (1981), *J. Phys. Soc. Jpn.* **55**, 3305 (1986), Y. Ozeki and H. Nishimori, *J. Phys. A: Math. Gen.* **26**, 3399 (1993).
- [8] Different variants of the phase diagram Fig. 1 have been presented before, in particular in [9] which we refer to for further comments and references regarding global properties of the phase diagram.
- [9] I. A. Gruzberg, N. Read and A. W. W. Ludwig, *Phys. Rev. B* **63**, 104422 (2001).
- [10] A. Georges and P. Le Doussal, *unpublished preprint* (1988), P. Le Doussal and A. B. Harris, *Phys. Rev. Lett.* **61**, 625 (1988), *Phys. Rev. B* **40**, 9249 (1989).
- [11] I. Morgenstern and K. Binder, *Phys. Rev. B* **22**, 288 (1980), I. Morgenstern and Horner, *Phys. Rev. B* **25**, 504 (1982), W. L. McMillan, *Phys. Rev. B* **28**, 5216 (1983).
- [12] A. Honecker, M. Picco and P. Pujol, *in preparation*.
- [13] W. L. McMillan, *Phys. Rev. B* **29**, 4026 (1984).
- [14] Y. Ozeki and H. Nishimori, *J. Phys. Soc. Jpn.* **56**, 3265 (1987).
- [15] R. R. P. Singh and J. Adler, *Phys. Rev. B* **54**, 364 (1996).
- [16] Y. Ozeki and N. Ito, *J. Phys. A: Math. Gen.* **31**, 5451 (1998).
- [17] F. D. A. Araújo Reis, S. L. A. de Queiroz and R. R. dos Santos, *Phys. Rev. B* **60**, 6740 (1999).
- [18] We omit a factor β since it is not important at finite temperature.
- [19] M. P. Nightingale, pp. 287-351 in: V. Privman (ed.), *Finite Size Scaling and Numerical Simulations of Statistical Physics*, World Scientific, Singapore (1990).
- [20] D. Stauffer and A. Aharony, *Introduction to Percolation Theory*, 2nd edition, Taylor & Francis, London (1994).
- [21] H. W. J. Blöte, J. L. Cardy and M. P. Nightingale, *Phys. Rev. Lett.* **56**, 742 (1986), I. Affleck, *Phys. Rev. Lett.* **56**, 746 (1986).